

GABA METABOLISM IN VENEZUELAN EQUINE  
ENCEPHALOMYELITIS VIRUS INFECTION.  
PRELIMINARY COMMUNICATION

Ernesto Bonilla, Slavia Ryder y Jesús Estevez\*

ABSTRACT

Mice infected with Venezuelan equine encephalomyelitis virus showed a significant decrease in GABA content of cerebral hemispheres. Activity of the enzyme which synthesizes GABA, glutamic acid decarboxylase, is also reduced in whole cerebral hemispheres, caudate nucleus, and frontal cortex of infected animals, as compared to values obtained from the same regions of control mice.

INTRODUCTION

Some workers believe that viral encephalitis is the major cause of Parkinson's disease (3). This disease, in turn, is characterized by a degeneration of the nigrostriatal dopaminergic pathway with the consequent depletion of dopamine and homovanillic acid in the substantia nigra and striatum (4, 6). A consistent decrease in glutamic acid decarboxylase (GAD) activity in the substantia nigra, caudate nucleus, putamen, and globus pallidus with no corresponding change in gamma-aminobutyric acid (GABA) concentration has also been found in post-mortem brain of Parkinsonian patients (8).

---

\* Instituto de Investigación Clínica, Apartado 1151, Facultad de Medicina, Universidad del Zulia. Maracaibo, Venezuela.

Recent reports suggest that viral infections can produce changes in the turnover of brain monoamines. These changes are characterized by increased concentrations of the acid end-products, homovanillic acid (HVA) and 5-hydroxyindoleacetic acid (9-11). A rise in dopamine (DA), HVA, and serotonin in Venezuelan equine encephalomyelitis (VEE) virus-infected brains has also been reported (1, 2).

The present work was undertaken in order to investigate the functional state of the GABA-ergic neurons in VEE virus infection. We now report studies which indicate that in this experimentally induced encephalitis there is a dysfunction of brain GABA metabolism.

## MATERIAL AND METHODS

Swiss albino mice, 21-28 days old, were inoculated intraperitoneally with 0.03 ml (containing 100 LD50) of the Guajira strain of VEE virus (14) suspended in 0.75% bovine albumin phosphate-buffered saline solution (BAPS), pH 7.4 (7). Control animals were injected with 0.03 ml of BAPS and killed simultaneously with the diseased animals. On the sixth day after inoculation, when paralysis of the limbs was evident, the mice were sacrificed by cervical dislocation, the brains immediately removed and analysed for GABA content (5) and GAD activity (12). None of the animals survived the infection after the seventh day of inoculation.

## RESULTS AND DISCUSSION

The data shown on table I demonstrated a significant decrease in GABA concentration in cerebral hemispheres of mice infected with VEE virus. A decrease in GAD activity was observed in whole cerebral hemispheres, frontal cortex and caudate nucleus (Table II). As shown in the present report the changes in GABA function seen in VEE virus infection are not identical to those reported in Parkinson's disease. As stated, in the latter disease there is a reduction in the activity of GAD in the striatum, but concentrations are normal (8).

Viral infections and the dysfunction of monoamine metabolism have both been considered to be responsible for the development of psychiatric and neurological diseases. In our laboratory we have noted that mice infected experimentally with VEE virus showed excitation and hypermotility followed by hypomotility, paralysis, coma, and death. Episodes of emotional and conductual disturbances as well as psychotic behaviour are common in human cases of VEE viral infections (13).

### Acknowledgements

We thank Mario Salazar for technical assistance. This work was supported by a grant from CONICIT (51.26.S1.0870).

**TABLE I**

GABA CONTENT IN CEREBRAL HEMISPHERES OF MICE INFECTED WITH VEE VIRUS. DATA REPRESENT MEANS  $\pm$  S.E. OF 8 DUPLICATED EXPERIMENTS.

Groups	GABA concentration (n moles/mg wet tissue)
Control	1.75 $\pm$ 0.15
Infected	0.84 $\pm$ 0.10
P	< 0.001

**TABLE II**

ACTIVITY OF L-GLUTAMIC ACID DECARBOXYLASE IN DISCRETE BRAIN REGIONS OF CONTROL AND VEE VIRUS INFECTED MICE. ENZYMATIC ACTIVITY EXPRESSED AS N MOLES CO<sub>2</sub> MIN/G WET WT. RESULTS ARE GIVEN AS MEANS  $\pm$  S.E. OF DUPLICATED OBSERVATIONS. FIGURES IN PARENTHESES REPRESENT THE NUMBER OF EXPERIMENTS.

Groups	Cerebral hemispheres	Caudate nucleus	Frontal cortex
Control	441.61 $\pm$ 26.11 (6)	544.22 $\pm$ 37.26 (8)	435.88 $\pm$ 30.16 (8)
Infected	318.88 $\pm$ 37.40 (6)	315.65 $\pm$ 30.16 (8)	274.84 $\pm$ 12.78 (7)
P	< 0.01	< 0.0001	< 0.0001

## RESUMEN

**Metabolismo de GABA en la infección producida por el virus de la encefalitis equina venezolana.** Bonilla E, Ryder S, Estevez J (Instituto de Investigación Clínica, Apartado 1151, Maracaibo, Venezuela). *Invest Clín* 19 (1): 41-45, 1978.- En ratones infectados con el virus de la encefalomyelitis equina venezolana, se observó un descenso significativo en la concentración de GABA en los hemisferios cerebrales. La actividad de la enzima responsable de la síntesis de GABA (decarboxilasa del ácido glutámico) también se encuentra reducida en los hemisferios cerebrales, núcleo caudado y corteza frontal de los animales infectados.

## REFERENCES

- 1- BONILLA E, RYDER S: Venezuelan equine encephalomyelitis virus infection: effect on dopamine metabolism of mouse brain. Preliminary communication. *Invest Clín* 14(2): 82-86, 1973.
- 2- BONILLA E, RYDER S, HERNANDEZ H: Venezuelan equine encephalomyelitis virus infection: effect on monoamine metabolism of mouse brain. *J Neurochem* 25(4): 529-530, 1975.
- 3- CALNE DB: In: Parkinsonism, physiology, pharmacology and treatment. Arnold E, ed., London, 1970, p 18.
- 4- EHRINGER H, HORNYKIEWICZ O: Verteilung von Noradrenalin und Dopamin (3-Hydroxytyramin) im Gehirn des Menschen und ihr Verhalten bei Erkrankungen des extrapyramidalen system. *Klin Wochschr* 38: 1236-1239, 1960.
- 5- FAHN S, COTE LJ: Regional distribution of gamma-aminobutyric acid (GABA) in brain of the Rhesus monkey. *J Neurochem* 15: 209-213, 1968.
- 6- FAHN S, LIBSCH LR, CUTLER RW: Monoamines in the human neostriatum: topographic distribution in normals and in Parkinson's disease and their role in akinesia, rigidity, chorea, and tremor. *J Neurol Sci* 14: 427-455, 1971.
- 7- HAMMON W, SATHER GE: In: Diagnostic procedures for viral and rickettsial infections. Lennette EH, Schmidt NJ, eds. Amer Public Hlth Ass, New York, 1969, pp 227-280.

- 8- HORNYKIEWICZ O, LLOYD KG, DAVIDSON L: In: GABA in nervous system function. Roberts E, Chase TN, Towers DB, eds. Raven Press, New York, 1976, p 479.
  - 9- LYCKE E, MODIGH K, ROOS BE: The monoamine metabolism in viral encephalitides of the mouse. Part 1: Virological and biochemical results. *Brain Res* 23: 235-246, 1970.
  - 10- LYCKE E, ROOS BE: The monoamine metabolism in viral encephalitides of the mouse. Part 2: Turnover of monoamines in mice infected with herpes simplex virus. *Brain Res* 44: 603-613, 1972.
  - 11- LYCKE E, ROOS BE: Virus infection in infant mice causing persistent impairment of turnover of brain catecholamines. *J Neurol Sci* 26: 49-60, 1975.
  - 12- MacDONNELL PC, GREENGARD O: The distribution of glutamate decarboxylase in rat tissues; isotopic vs fluorimetric assays. *J Neurochem* 24: 615-618, 1975.
  - 13- NEGRETTE A, MOSQUERA J: Epidemia de encefalitis de 1959 en Maracaibo (San Francisco), Estado Zulia, Venezuela. Manifestaciones clínicas y terapéutica antibiótica. *Invest Clín* 15(4): 11-44, 1974.
  - 14- SOTO ESCALONA A, FINOL LT, RYDER S: Estudio de un brote de encefalitis venezolana en el Distrito Páez, Estado Zulia, en octubre de 1968. *Invest Clín* 10(31): 45-57, 1969.
-