

---

---

# Gender-based comparative analysis of knee injury risk during cutting maneuvers in non-professional athletes: a kinetic and kinematic perspective.

Adrián Feria-Madueño<sup>1</sup>, Timothy E. Hewett<sup>2,3</sup> and Borja Sañudo<sup>1</sup>

<sup>1</sup> Department of Physical Education and Sport, University of Seville. Pirotecnia Street, Seville, Spain.

<sup>2</sup> Department of Orthopedics, Marshall University Biomechanics Laboratories and Sports Medicine Research, Marshall University, Huntington, West Virginia, USA.

<sup>3</sup> Hewett Global Consulting, Newport, KY.

**Keywords:** anterior cruciate ligament; Q angle; accelerometry; force platform.

**Abstract.** This study aimed to compare the risk of knee injury between men and women by integrating kinetic and kinematic parameters in a non-professional athlete population. Two hundred non-professional athletes were recruited for the present study. Three change of direction tests were conducted, consisting of two open cuts at 30 (SC<sub>30</sub>) and 45 degrees (SC<sub>45</sub>) and one closed cut at 45 degrees (SC<sub>45cl</sub>). Kinetic variables, including three-dimensional force and accelerations in the three axes of movement and ground contact time, were assessed using force platform and accelerometers. The initial and maximum angles of the ankle, knee, hip, and trunk were analyzed by photogrammetry. The data was compared between males and females to examine gender differences. Gender analysis demonstrated significant differences in force values, with women displaying higher medial-lateral (ML) force in SC<sub>30</sub> and men exhibiting higher vertical ground reaction force (VGRF) and anterior-posterior (AP) force in SC<sub>45</sub>. Gender-specific analysis indicated higher partial knee accelerations in women during SC<sub>30</sub> and SC<sub>45</sub>, with significant differences observed in acceleration in the vertical axis. Gender differences were observed in certain kinematic variables, with women displaying higher ankle flexion at initial contact in SC<sub>30</sub> and higher ankle flexion at maximum flexion and ankle dorsiflexion range in SC<sub>45</sub>. Men showed lower knee flexion angles in both SC<sub>45cl</sub> and SC<sub>45</sub>. These findings provide valuable insights into the kinetics and kinematics of change of direction movements and highlight gender-specific differences that may have implications for training and injury prevention strategies. Further research is needed to understand the underlying factors contributing to these differences and their impact on performance and injury risk.

## **Análisis comparativo, basado en el género, del riesgo de lesión de rodilla durante cambios de dirección en atletas no profesionales: una perspectiva cinética y cinemática.**

*Invest Clin 2024; 65 (2): 192 – 205*

**Palabras clave:** ligamento cruzado anterior; ángulo Q; acelerómetro; plataforma de fuerza.

**Resumen.** El objetivo de este estudio fue comparar el riesgo de lesión de rodilla entre hombres y mujeres mediante la integración de parámetros cinéticos y cinemáticos en una población de atletas no profesionales. Doscientos atletas no profesionales fueron reclutados para el presente estudio. Se realizaron tres pruebas de cambio de dirección, consistentes en dos cortes abiertos a 30 (SC<sub>30</sub>) y 45 grados (SC<sub>45</sub>), y un corte cerrado a 45 grados (SC<sub>45cl</sub>). Se evaluaron las variables cinéticas, incluida la fuerza tridimensional, así como las aceleraciones en los tres ejes de movimiento y el tiempo de contacto con el suelo mediante plataforma de fuerza y acelerometría. Se analizaron los ángulos iniciales y máximos del tobillo, la rodilla, la cadera y el tronco a través de fotogrametría. Se realizó una comparación de los datos entre hombres y mujeres para examinar las diferencias de género. El análisis de género demostró diferencias significativas en los valores de fuerza, mostrando las mujeres una mayor fuerza medial-lateral (ML) en SC<sub>30</sub>, y los hombres una mayor fuerza de reacción vertical al suelo (VGRF) y fuerza anteroposterior (AP) en SC<sub>45</sub>. Los análisis específicos de género indicaron mayores aceleraciones parciales de rodilla en las mujeres durante SC<sub>30</sub> y SC<sub>45</sub>, observándose diferencias significativas en la aceleración en el eje vertical. Se observaron diferencias de género en determinadas variables cinemáticas, mostrando las mujeres mayor flexión del tobillo en el contacto inicial en SC<sub>30</sub> y mayor flexión del tobillo en flexión máxima y rango de dorsiflexión del tobillo en SC<sub>45</sub>. Los hombres mostraron ángulos de flexión de rodilla más bajos tanto en SC<sub>45cl</sub> como en SC<sub>45</sub>. Estos resultados proporcionan información valiosa sobre la cinética y la cinemática de los movimientos de cambio de dirección y ponen de relieve las diferencias específicas de género que pueden tener implicaciones para las estrategias de entrenamiento y prevención de lesiones. Se necesita más investigación para comprender mejor los factores subyacentes que contribuyen a estas diferencias y su impacto en el rendimiento y el riesgo de lesiones.

*Received: 12-10-2023*

*Accepted: 08-02-2024*

### **INTRODUCTION**

Gender is a significant factor to consider in the incidence and prevention of knee injuries during changes of direction<sup>1,2</sup>. One of the structures with a higher incidence of injury is the anterior cruciate ligament

(ACL), and research has demonstrated that women have an increased risk compared to men in sports involving frequent changes of direction and deceleration<sup>3</sup>.

From a kinetic standpoint, the mechanism of injury during knee changes of directions involves a combination of torsional and

axial loading forces applied to the ACL<sup>3</sup>. The knee is vulnerable during direction changes due to the combination of rotational and axial loading movements on the ligament<sup>4</sup>. The assessment of the ACL using force platforms is well-established<sup>5-7</sup>. However, there is limited evidence on its analysis using other parameters such as accelerometry in each axis and ground contact time during changes of directions.

In women, hip anatomy and knee biomechanics during changes of directions may contribute to a higher incidence of ACL injuries<sup>8</sup>. Women have wider hips and a larger Q angle, which can increase internal knee rotation during direction changes and thus increase the risk of ACL injuries<sup>4</sup>. Consequently, women exhibit a smaller flexion angle and greater dynamic valgus during knee changes, associated with lower knee stability and an increased risk of injury<sup>3</sup>.

These kinetic and kinematic factors related to the risk of knee injuries include not only the direction and magnitude of forces applied to the knee during changes of direction<sup>9</sup>, but also depend on the knee angle maintained and ground contact time during changes of direction<sup>10</sup>. However, there is limited research integrating both factors. Despite knowledge of the risk factors most associated with an increased risk of knee injuries, there is a lack of evidence in evaluating these risk factors in a non-professional

athlete population, necessitating studies that assess kinetic and kinematic parameters, integrating them into a comprehensive analysis with innovative parameters such as acceleration or contact time.

Therefore, this study aimed to compare the risk of knee injury between men and women by integrating kinetic and kinematic parameters in a non-professional athlete population.

## MATERIALS AND METHODS

### Subjects

Two hundred non-professional athletes were recruited for the present study. After the initial meeting, the inclusion criteria were explained. These criteria consisted of engaging in physical activity for at least 30 minutes per day, three days per week, and having no history of knee or ankle injury in the past 12 months. Following this meeting, 38 athletes withdrew from the study, resulting in one hundred sixty-two participants (75% males, 25% females). All participants completed surveys to assess their weekly and daily levels of physical activity, as well as their height and body mass (Table 1). The potential risks of the tests were verbally communicated to the participants, and they provided signed informed consent. The Ethics Committee of the University of Seville approved the study.

**Table 1**  
Characteristics of the participants.

Variables	Mean	Men (n=122)	Women (n=40)	<i>p</i>
Age (years)	24 ± 3	25 ± 2	22 ± 1	0.110
Body mass (Kg)	72.84 ± 12.76	76.92 ± 8.12	65.36 ± 4.52	0.103
Height (m)	1.74 ± 0.07	1.80 ± 0.09	1.63 ± 0.06	0.071
BMI (kg/m <sup>2</sup> )	23.78 ± 2.86	24.75 ± 1.12	21.34 ± 2.12	0.096
PA (hours/week)	8.38 ± 4.01	8.79 ± 3.96	7.91 ± 2.34	0.101
PA <sub>day</sub> (hours/day)	1.97 ± 1.66	2.03 ± .098	1.86 ± 1.10	0.510

\**p* ≤ 0.05 Men vs Women. Statistical analysis was done using a repeated-measures two-way ANOVA. Results expressed as mean ± standard error of the mean. BMI: body mass index. PA: physical activity.

## Procedures

The participants completed a Maximum Voluntary Contraction (MVC) test using the Biodex Multi-joint System (Shirley, New York). The exercise was knee flexion at 45° to assess the strength of the posterior thigh musculature and knee extension at 45° to assess the anterior thigh musculature. Following this, they underwent a directional change test with the dominant foot, which included two open cuts at 45° (SC<sub>45</sub>) and 30° (SC<sub>30</sub>), as well as a closed cut (SC<sub>45cl</sub>) on a force platform (Kistler 9260 AA6, Winterthur, Switzerland). Prior to the test, all participants underwent a standardized warm-up, which involved a 5-minute cycling session on a cycloergometer (Ergoline 900, Ergometrics, Bitz, Germany) at an intensity of 60 W (60 rpm). Additionally, they familiarized themselves with sports-specific side-cutting maneuvers through five to eight practice attempts. The velocity of the movements was regulated using a metronome set at 4-5.5 m/s, and the designated direction for the movements was indicated on the floor using tape.

## Kinetics

A force plate (Kistler 9260 AA6, Winterthur, Switzerland) assessed the Ground Reaction Forces (GRF) in the vertical, mediolateral, and anteroposterior axes. Additionally, triaxial accelerometers (xyz-PLUX, PLUX-Wireless Biosignals, S.A., Lisbon, Portugal) were employed to measure accelerations at the knee and ankle joints. The knee accelerometer was positioned on the lateral condyle. In contrast, the ankle accelerometer was placed on the malleolus (Fig. 1). Antero-posterior (AP), Medial-Lateral (ML), and Longitudinal (Z) axes were calculated, and all signals were recorded at a sampling rate of 1000 Hz. Contact Time was determined during side-cutting by measuring the duration from the initial ground contact until the foot entered the flight phase.



Fig. 1. Location of accelerometers.

## Kinematics

An analysis of knee, ankle, hip, and trunk angles was conducted. Reflective markers were positioned on the lateral malleolus, lateral condyle, greater trochanter, and acromion to capture the movements precisely. The angles were calculated at two specific time points: first, during the initial contact between the foot and the ground, and second, when the Vertical Ground Reaction Force (VGRF) was recorded. Three video cameras (240 Fps) were positioned perpendicularly, one for each type of change of direction, at a distance of two meters from the central point of the force platform. Subsequently, the Kinovea software was employed for digitization and angle extraction by 2D analysis.

## Statistical analysis

Statistical analysis was performed using the SPSS 22.0 software package (SPSS Inc., Chicago, IL, USA). The Kolmogorov-Smirnov

test assessed the data distribution, and parametric variables were identified. Baseline data were compared using an independent-sample t-test. Potential statistical differences were evaluated using a repeated-measures two-way ANOVA (time x group). Mean values are reported with standard deviation (SD), and statistical significance was set at  $p \leq 0.05$ .

The effect size (Cohen's *d*) was calculated by dividing the difference between the means of the groups by the combined standard deviation of both groups, considering the sample size. Effect sizes of 0.2 were considered small, 0.5 moderate, and 0.8 large.

## RESULTS

### Kinetic

During the SC<sub>30</sub>, subjects exhibited a higher value of VGRF of  $2135.49 \pm 633.14$  N, which corresponds to approximately three times their body weight, while ML Force ( $625.44 \pm 170.99$  N) and AP Force ( $945.62 \pm 421.53$  N) were lower. For the SC<sub>45</sub>, VGRF was lower compared to SC<sub>30</sub>, with a value of  $2015.83 (\pm 687.27)$  N, while ML Force was  $566.87 (\pm 189.22)$  N and AP Force was  $765.75 (\pm 417.06)$  N. In SC<sub>45cl</sub>, VGRF was

$1790.60 (\pm 517.55)$  N, and ML Force exhibited the smallest force value among all measured changes of directions ( $515.41 \pm 181.53$  N), as well as in AP Force ( $670.37 \pm 368.42$  N). For more specific results, the force was normalized to each subject's body weight for each directional change, as shown in Table 2.

Regarding the SC<sub>30</sub>, the results showed the highest applied force in all axes, including VGRF, AP Force, and ML Force. The gender analysis in Table 2 revealed that during the SC<sub>30</sub> directional change, women exhibited significantly higher force values than men in ML Force ( $p = 0.045$ ). On the other hand, in the SC<sub>45</sub>, men demonstrated higher values than women in both VGRF ( $p = 0.025$ ) and AP Force ( $p = 0.020$ ). Finally, in SC<sub>45cl</sub>, significant differences were only observed in the force exerted in AP Force, where women displayed a higher force value than males ( $p = 0.020$ ).

The analysis of partial knee and ankle accelerations during changes of directions is presented in Table 3. The SC<sub>30</sub> exhibited the lowest ACC<sub>KNEE</sub>AP ( $1.72 \pm 1.48$  g). Similarly, ACC<sub>ANKLE</sub>Z had the lowest values compared to the other changes of directions ( $4.11 \pm$

**Table 2**  
Descriptive analysis of force during direction change.

	Variables	Mean	Men	Women	<i>p</i>	d-Cohen
SC <sub>30</sub>	VGRF (N/Kg)	29.49±7.75	29.54±7.42	29.33±8.81	0.465	0.02
	ForceAP (N/Kg)	12.99±5.32	13.54±5.45	11.26±4.53	0.159	0.43
	ForceML (N/Kg)	8.64±2.08	8.60±1.83	8.75±2.76	0.045*	-0.07
SC <sub>45</sub>	VGRF (N/Kg)	27.70±7.81	28.20±8.32	26.14±5.76	0.025*	0.26
	ForceAP (N/Kg)	10.54±5.39	11.14±5.68	8.68±3.90	0.020*	0.46
	ForceML (N/Kg)	7.78±2.09	7.96±2.15	7.23±1.77	0.568	0.35
SC <sub>45cl</sub>	VGRF (N/Kg)	25.13±7.59	24.02±6.92	28.60±8.61	0.159	-0.62
	ForceAP (N/Kg)	9.35±5.09	9.02±4.54	10.39±6.48	0.020*	-0.26
	ForceML (N/Kg)	7.17±2.40	7.13±2.40	7.32±2.45	0.612	-0.07

\* $p \leq 0.05$  Men vs Women. Statistical analysis was done using a repeated-measures two-way ANOVA. Results expressed as mean±standard error of the mean. d-Cohen = the effect size. VGRF (N) = Ground reaction force. ForceML (N) = Force in medio-lateral axe. ForceAP (N) = Force in antero-posterior axe.

**Table 3**  
Descriptive analysis of the acceleration in the three axes during direction changes.

	SC <sub>30</sub>	SC <sub>45</sub>	SC <sub>45cl</sub>
ACC <sub>KNEE_ML</sub>	2.46±1.50	2.39±1.58	2.49±1.49
ACC <sub>KNEE_AP</sub>	1.72±1.48	2.03±1.55	2.08±1.55
ACC <sub>KNEE_Z</sub>	1.63±1.37	1.53±1.30	1.69±1.40
ACC <sub>ANKLE_ML</sub>	4.19±0.78	4.23±0.75	4.14±0.88
ACC <sub>ANKLE_AP</sub>	4.04±0.98	4.05±0.92	3.98±0.99
ACC <sub>ANKLE_Z</sub>	4.11±0.81	4.13±0.68	4.20±0.79

Results expressed as mean ± standard error of the mean. ACC<sub>KNEE\_ML</sub> (g) = Acceleration in medio-lateral axis in knee. ACC<sub>KNEE\_AP</sub> (g) = Acceleration in antero-posterior axis in knee. ACC<sub>KNEE\_Z</sub> (g) = Acceleration in vertical axis in knee. ACC<sub>ANKLE\_ML</sub> (g) = Acceleration in medio-lateral axis in ankle. ACC<sub>ANKLE\_AP</sub> (g) = Acceleration in antero-posterior axis in ankle. ACC<sub>ANKLE\_Z</sub> (g) = Acceleration in vertical axis in ankle.

0.81). For the SC<sub>45</sub>, both ACC<sub>KNEE\_ML</sub> (2.39 ± 1.58 g) and ACC<sub>KNEE\_Z</sub> (1.53 ± 1.30 g) showed the lowest values. However, both ACC<sub>ANKLE\_ML</sub> and ACC<sub>ANKLE\_AP</sub> were higher than in the other changes of directions (4.23 ± 0.75 g and 4.05 ± 0.92 g, respectively). During the SC<sub>45cl</sub>, the highest ACC<sub>KNEE\_ML</sub> (2.49 ± 1.49 g), ACC<sub>KNEE\_Z</sub> (1.69 ± 1.40 g), and ACC<sub>ANKLE\_Z</sub> (4.20 ± 0.79 g) were found. Conversely, ACC<sub>ANKLE\_ML</sub> and ACC<sub>ANKLE\_AP</sub> were lower during this directional change (4.14 ± 0.88 g and 3.98 ± 0.99 g, respectively).

On the other hand, gender-specific acceleration was also evaluated (Table 4). During the SC<sub>30</sub>, although no significant differences were observed, partial knee accelerations were higher in women than men. In the ankle, only ACC<sub>ANKLE\_Z</sub> showed higher values in women than men (4.20 ± 0.76 g). In the case of SC<sub>45</sub>, there was a trend towards higher acceleration in women compared to men, both in the knee and ankle. Regarding ACC<sub>KNEE\_Z</sub>, the difference between women and men was significant (p=0.002), with women displaying an acceleration of 1.82 ± 1.54 g compared to 1.44 ± 1.21 g exhibited by men.

During the SC<sub>45cl</sub>, no significant gender differences were found, except for ACC<sub>KNEE\_ML</sub> where men exhibited higher values (2.49 ± 1.48 g). In the remaining partial knee accelerations, women showed higher

values for this parameter, although they were not statistically significant. Regarding the ankle, only a trend towards higher values in men than women was observed for ACC<sub>ANKLE\_Z</sub> (4.22 ± 0.81 g), although statistical significance was not reached.

### Kinematics

Kinematic factors were also studied during changes of direction. Knee flexion angles at initial contact (KneeFlexInit) and maximum knee flexion angle (KneeFlexMax) were analyzed. The range of ankle dorsiflexion (AnkleDorsiRange), initial hip flexion angle (HipFlexInit), maximum hip flexion angle (HipFlexMax), initial trunk flexion angle (TrunkFlexInit), and maximum trunk flexion angle (TrunkFlexMax) were also analyzed.

Regarding the changes of directions (Table 5), the highest value of KneeFlexInit was observed in SC<sub>30</sub> (144.6° ± 7.6°), while the lowest value of KneeFlexMax was found during SC<sub>45</sub> (125.2° ± 8.9°). For the hip, the maximum value of HipFlexMax was 60.6° during SC<sub>30</sub>, while the trunk flexed 92.9°. Finally, a range of ankle dorsiflexion of 39.4° was found between the initial contact of the directional change and the maximum plantar flexion during the change.

In SC<sub>45</sub>, KneeFlexMax was 125.2° (extension), similar to the value of 123.3° in

**Table 4**  
Differences between men and women in acceleration during direction changes.

	Variables	Men	Women	<i>p</i>	d-Cohen
SC <sub>30</sub>	ACC <sub>KNEE_ML</sub>	2.33±1.47	2.86±1.56	0.558	-0.35
	ACC <sub>KNEE_AP</sub>	1.66±1.50	1.91±1.44	0.573	-0.16
	ACC <sub>KNEE_Z</sub>	1.57±1.30	1.84±1.55	0.080	-0.19
	ACC <sub>ANKLE_ML</sub>	4.20±0.73	4.14±0.94	0.215	0.07
	ACC <sub>ANKLE_AP</sub>	4.04±0.96	4.02±1.06	0.433	0.02
	ACC <sub>ANKLE_Z</sub>	4.08±0.83	4.20±0.76	0.413	-0.14
SC <sub>45</sub>	ACC <sub>KNEE_ML</sub>	2.25±1.54	2.81±1.66	0.516	-0.35
	ACC <sub>KNEE_AP</sub>	1.92±1.57	2.35±1.49	0.563	-0.27
	ACC <sub>KNEE_Z</sub>	1.44±1.21	1.82±1.54	0.002*	-0.29
	ACC <sub>ANKLE_ML</sub>	4.20±0.71	4.32±0.86	0.890	-0.16
	ACC <sub>ANKLE_AP</sub>	4.04±0.92	4.09±0.95	0.483	-0.05
	ACC <sub>ANKLE_Z</sub>	4.09±0.70	4.27±0.60	0.344	-0.26
SC <sub>45ml</sub>	ACC <sub>KNEE_ML</sub>	2.49±1.48	2.48±1.55	0.866	0.01
	ACC <sub>KNEE_AP</sub>	1.96±1.53	2.46±1.58	0.795	-0.32
	ACC <sub>KNEE_Z</sub>	1.64±1.37	1.85±1.51	0.200	-0.14
	ACC <sub>ANKLE_ML</sub>	4.11±0.87	4.22±0.90	0.937	-0.12
	ACC <sub>ANKLE_AP</sub>	3.97±1.01	4.01±0.96	0.965	-0.04
	ACC <sub>ANKLE_Z</sub>	4.22±0.81	4.15±0.71	0.900	0.08

\**p* ≤ 0.05 Men vs Women. Statistical analysis was done using a repeated-measures two-way ANOVA. Results expressed as mean±standard error of the mean. d-Cohen = the effect size. ACC<sub>KNEE\_ML</sub> (g) = Acceleration in medio-lateral axe in knee. ACC<sub>ANKLE\_AP</sub> (g) = Acceleration in antero-posterior axe in knee. ACC<sub>ANKLE\_Z</sub> (g) = Acceleration in vertical axe in knee. ACC<sub>ANKLE\_ML</sub> (g) = Acceleration in medio-lateral axe in ankle. ACC<sub>ANKLE\_AP</sub> (g) = Acceleration in antero-posterior axe in ankle. ACC<sub>ANKLE\_Z</sub> (g) = Acceleration in vertical axe in ankle.

**Table 5**  
Description of the kinematics in the experimental phase in direction change.

Variable	SC <sub>30</sub>	SC <sub>45</sub>	SC <sub>45cl</sub>
	Mean	Mean	Mean
Angle <sub>in</sub> Knee (degrees °)	144.63±7.65	143.45±17.86	141.43±9.87
Angle <sub>max</sub> Knee (degrees °)	123.33±10.00	125.16±8.91	119.97±14.18
Angle <sub>in</sub> Hip (degrees °)	42.50±2.94	41.54±5.63	40.92±4.56
Angle <sub>max</sub> Hip (degrees °)	60.57±5.50	51.21±6.36	54.37±5.37
Angle <sub>in</sub> Trunk (degrees °)	104.54±6.35	105.15±5.15	98.98±4.76
Angle <sub>max</sub> Trunk (degrees °)	92.99±5.36	95.02±5.73	77.77±6.54
Angle <sub>in</sub> Ankle (degrees °)	99.12±6.87	98.92±5.30	98.82±5.25
Angle <sub>max</sub> Ankle (degrees °)	59.70±7.45	57.19±5.40	56.97±4.28
Ankle_Dorsi_Angle (degrees °)	39.42±3.32	41.72±5.38	41.84±4.92

Results expressed as mean±standard error of the mean. Angle<sub>in</sub> Knee = Angle of knee in the first contact. Angle<sub>max</sub> Knee = Angle of knee in the maximum flexion. Angle<sub>in</sub> Hip = Angle of hip in the first contact. Angle<sub>max</sub> Hip = Angle of hip in the maximum flexion. Angle<sub>in</sub> Trunk = Angle of trunk in the first contact. Angle<sub>max</sub> Trunk = Angle of trunk in the maximum flexion. Angle<sub>in</sub> Ankle = Angle of ankle in the first contact. Angle<sub>max</sub> Ankle = Angle of ankle in the maximum flexion. Ankle\_Dorsi\_Angle = Range of ankle dorsiflexion.

SC<sub>30</sub>. At that moment, HipFlexMax was 51.21°, while TrunkFlexMax showed a flexion of 95.1°. Additionally, the AnkleDorsiRange was 41.7°, defined as the amount of flexion from the first contact until maximum flexion occurred.

In SC<sub>45cl</sub>, KneeFlexMax was 119.9°, representing the most significant flexion of the three change types. From initiating contact to maximum knee flexion, the hip flexed by 13.5°. As for the trunk, it exhibited a flexion of 77.8° at that moment. Finally, the AnkleDorsiRange was 41.8°.

Finally, a gender analysis of kinematics during direction changes was performed (Table 6). It was observed that during SC<sub>30</sub>, women exhibited a significantly higher AnkleFlexInit than men ( $p=0.040$ ). However, gender differences during SC<sub>45</sub> were found in AnkleFlexMax ( $p=0.004$ ) and AnkleDorsiRange ( $p=0.007$ ). Lastly, during SC<sub>45cl</sub>, gender differences were found in KneeFlexInit ( $p<0.001$ ) and KneeFlexMax ( $p=0.006$ ), where men displayed a lower angle compared to women.

## DISCUSSION

The main objective of this study was to compare the risk of knee injury between men and women through the analysis of kinetic and kinematic parameters in a population of non-professional athletes. Both kinetic aspects, such as GRF and accelerations, as well as kinematic elements, such as angles at the moment of directional change, have proven to be crucial, with women likely presenting a higher risk of injury.

### Kinetics

During changes of directions, the VGRF relative to body weight was significantly higher in SC<sub>30</sub> (29.49 N/kg  $\pm$  7.75) compared to SC<sub>45</sub> (27.70 N/kg  $\pm$  7.81;  $p=0.004$ ) and SC<sub>45cl</sub> (25.13 N/kg  $\pm$  7.59;  $p<0.001$ ). One possible explanation for these differences is that greater force needs to be exerted in the medial-lateral (ML) axis to perform

more forceful or larger amplitude changes (remember that 30° is measured above the horizontal plane). According to Nigg's paradigm<sup>11</sup>, which compiled studies spanning over 25 years, it became evident that VGRF plays a particularly important role in injuries among individuals engaged in physical activities such as running, especially in situations where pronation is accentuated, such as during changes of directions. This possibility is also supported by the values of ML force found in our study, as the analyzed subjects displayed significantly higher values during SC<sub>30</sub> (8.64 N/kg  $\pm$  2.08) compared to SC<sub>45</sub> (7.78 N/kg  $\pm$  2.08,  $p<0.001$ ) and SC<sub>45cl</sub> (7.17 N/kg  $\pm$  2.40,  $p<0.001$ ). Therefore, higher levels of force, especially in the ML axis, which need to be attenuated by the lower extremities, suggest the need for increased preventive strategies in these types of actions common in various sports. It is essential to highlight that these 30°-changes of directions do not only increase the ML force but also the AP force (12.9 N/kg  $\pm$  5.32) when compared to SC<sub>45</sub> (10.54 N/kg  $\pm$  5.39,  $p<0.001$ ) or SC<sub>45cl</sub> (9.35 N/kg  $\pm$  5.09,  $p<0.001$ ), suggesting that biomechanical modifications in the different involved joints occur in response to these increments, as will be discussed later.

Despite numerous studies regarding the biomechanical and kinetic factors manifested during direction changes, the results are contradictory<sup>12</sup>. The aspect that seems most relevant in explaining this disparity of results may be the variety of procedures carried out in different studies. Brughelli *et al.*<sup>13</sup> differentiate between anticipated and unanticipated changes of directions, changes towards the dominant or non-dominant foot, as elements that influence the final results. Regardless of the methodological variety, it is evident that both VGRF, ML-force, and AP-force are variables closely related to the onset of injury processes<sup>2</sup> and should be controlled and attenuated as much as possible.

While it is evident that higher forces in both axes are associated with a greater risk



**Table 6**  
Kinematics during direction changes and differences between men and women.

Variables	SC <sub>30</sub>		p	d-Cohen	SC <sub>45</sub>		p	d-Cohen	SC <sub>45cl</sub>		p	d-Cohen
	Men	Women			Men	Women			Men	Women		
Angle <sub>in</sub> Knee (degrees °)	144.04±7.48	146.56±8.06	0.080	-0.33	145.33±7.53	145.13±6.43	0.891	-0.23	139.88±9.39	146.32±10.01	0.000*	-0.67
Angle <sub>max</sub> Knee (degrees °)	122.86±9.28	125.13±11.97	0.229	-0.23	124.74±8.80	126.75±9.22	0.239	-0.09	118.20±14.34	125.56±12.40	0.006*	-0.53
Angle <sub>in</sub> Hip (degrees °)	42.63±3.00	42.13±2.77	0.366	0.17	41.42±5.66	41.94±5.68	0.623	-0.15	40.93±4.74	41.00±4.01	0.938	-0.02
Angle <sub>max</sub> Hip (degrees °)	60.97±5.28	59.37±6.12	0.125	0.29	51.01±6.67	51.97±5.29	0.427	-0.05	54.29±5.42	54.43±5.22	0.891	-0.03
Angle <sub>in</sub> Trunk (degrees °)	104.45±6.53	104.72±5.87	0.819	-0.04	105.07±5.39	105.32±4.39	0.799	-0.12	98.91±4.87	99.08±4.44	0.855	-0.04
Angle <sub>max</sub> Trunk (degrees °)	93.09±5.56	92.59±4.76	0.624	0.09	94.82±5.92	95.51±5.10	0.524	0.04	78.00±6.61	76.72±6.18	0.299	0.20
Angle <sub>in</sub> Ankle (degrees °)	98.47±6.58	101.13±7.55	0.040*	-0.39	99.01±5.51	98.81±4.56	0.837	0.56	98.59±4.98	99.51±6.12	0.357	-0.17
Angle <sub>max</sub> Ankle (degrees °)	59.10±7.22	61.48±8.05	0.091	-0.32	57.91±5.29	54.97±5.25	0.004*	-0.52	57.13±4.31	56.37±4.24	0.351	0.18
Ankle_Dorsi_Angle (degrees °)	39.36±3.26	39.64±3.56	0.658	-0.08	41.10±5.10	43.83±5.81	0.007*	-0.23	41.46±4.81	43.13±5.18	0.072	-0.34

\*p ≤ 0.05 Men vs Women. Statistical analysis was done using a repeated-measures two-way ANOVA. Results expressed as mean ± standard error of the mean. d-Cohen = the effect size. Angle<sub>in</sub> Knee = Angle of knee in the first contact. Angle<sub>max</sub> Knee = Angle of knee in the maximum flexion. Angle<sub>in</sub> Hip = Angle of hip in the first contact. Angle<sub>max</sub> Hip = Angle of hip in the maximum flexion. Angle<sub>in</sub> Trunk = Angle of trunk in the first contact. Angle<sub>max</sub> Trunk = Angle of trunk in the maximum flexion. Angle<sub>in</sub> Ankle = Angle of ankle in the first contact. Angle<sub>max</sub> Ankle = Angle of ankle in the maximum flexion. Ankle\_Dorsi\_Angle = Range of ankle dorsiflexion.

of injury in this population group, it is essential to mention the differences observed in the different variables according to gender. Sigward and Powers<sup>14</sup> suggested that an increased risk of knee injury through increased valgus can occur due to greater forces in either axis in both men and women. In fact, their work reflected values that were very similar to those found in the present study regarding VGRF during changes of directions. Our results show that during SC<sub>30</sub>, where women sustained higher relative loads than men ( $p=0.045$ ), there were significant differences in the force exerted in the ML axis. This finding has been widely observed in studies with athletes, where women have shown significantly greater valgus than men<sup>15</sup>. Differences in the ML axis between men and women could be related to an increased risk of injury in females due to the biomechanical position of the knee in abduction or adduction during ground contact<sup>1,16</sup>.

On the other hand, during SC<sub>45</sub>, the differences were found in VGRF ( $p=0.025$ ) and AP force ( $p=0.020$ ), where men significantly exceeded women. Our results are consistent with those of Liu *et al.*<sup>17</sup>, who related a reduction of VGRF and Force<sub>AP</sub> with a reduction of the load supported by the knee during open direction changes. However, authors such as James *et al.*<sup>18</sup> found the opposite during maximum speed changes of directions. One possible explanation for these differences could be the openness of the directional change at a non-maximal speed. SC<sub>45</sub> may involve less demand than SC<sub>30</sub>, so being a more vertical change and closer to straight-line running, it is possible that AP force is increased, as well as VGRF in men.

Another important finding was that women exhibited a significantly higher AP force during SC<sub>45cl</sub> ( $p=0.020$ ) than men. Our results align with those presented by McLean *et al.*<sup>2</sup>, who found higher AP force in women compared to men during a directional change with an angle of exit between 35° and 55° (women = 1.80BW [ $\pm$  0.54]; men 1.54BW [ $\pm$  0.76]). Moreover, although

not statistically significant, women in our study also showed higher VGRF during SC<sub>45cl</sub>. Recently, de Hoyo *et al.*<sup>19</sup> reported very similar data in male athletes who performed a closed change with a 60° opening (24.1 N/kg  $\pm$  8.4). Unfortunately, these authors only conducted their intervention with male subjects, making a comparison impossible. Nevertheless, there is limited research on the role of gender in VGRF during closed changes of directions. This is the first study to assess vertical force in men and women during closed changes.

What seems evident is the increased risk of knee injury through ML force values during changes of directions based on gender<sup>1</sup>. Authors like Sigward and Powers<sup>14</sup> indicated that women who exerted greater ML force also exhibited a greater valgus moment, which increased the risk of a knee injury. Although our results do not provide data on impulse outcomes, a possible contribution can be inferred from the partial knee and ankle acceleration. As described previously, women reported significantly higher ML force during SC<sub>30</sub> than men and a slightly higher trend was observed in this variable during SC<sub>45cl</sub>. In this regard, it can also be observed that women experience 18.53% more ML knee acceleration than men during SC<sub>30</sub> and 19.93% during SC<sub>45</sub>, with almost no difference (0.4%) during SC<sub>45cl</sub>. This ML knee acceleration could represent valgus moments, although this fact should be corroborated in future studies through a kinematic analysis in the frontal plane.

There were significant differences between the different changes of direction. ACC<sub>KNEE</sub>AP was significantly higher in SC<sub>45cl</sub> compared to SC<sub>30</sub> ( $p=0.006$ ). This finding suggests an increased risk in closed changes compared to open changes, as the load is mobilized more rapidly in the sagittal axis, potentially leading to an increase in tibial translation. However, ForceAP was significantly lower in SC<sub>45cl</sub> than SC<sub>30</sub> ( $p<0.001$ ). These differences also existed between open changes, as ACC<sub>KNEE</sub>AP was significantly

higher in SC<sub>45</sub> than SC<sub>30</sub> ( $p=0.023$ ). One possible explanation for these differences could be the biomechanical demand, with SC<sub>45</sub> being less demanding than SC<sub>30</sub>, resulting in a faster strategy during less open changes. Analyzing ACC<sub>KNEE</sub>ML between changes of directions, SC<sub>30</sub> showed a significantly higher acceleration than SC<sub>45</sub> ( $p<0.001$ ) and SC<sub>45cl</sub> ( $p<0.001$ ). Sigward and Powers<sup>14</sup> suggest that frontal plane control is necessary to reduce the relative risk of knee injury. Our data indicate that SC<sub>30</sub> poses an increased risk compared to the other changes of directions, characterized by higher ForceML and ACC<sub>KNEE</sub>ML, which are associated with an increase in knee valgus.

Regarding gender differences during changes of directions, significant differences were found only in SC<sub>45</sub> ( $p\leq 0.05$ ), with women exhibiting higher ACC<sub>KNEE</sub>Z than men. One possible explanation could be that Chapell *et al.*<sup>20</sup> suggested that women experience greater shear force in the tibia during ground contact. This negative strategy indicates that energy absorption by women is less efficient than in men, or at least faster, which may be related to an increased risk of knee injury. Surprisingly, this trend is not observed in SC<sub>30</sub>, which would be expected to be more demanding at this level. However, the trend still favored women, who presented 14.67% more ACC<sub>KNEE</sub>Z during SC<sub>30</sub> and 11.35% more during SC<sub>45cl</sub>.

### Kinematics

It is evident that a knee angle close to the extension increases the risk of injury<sup>20-22</sup>. Our results indicate a higher KneeAnglein during SC<sub>30</sub> compared to the other changes of directions ( $144.63^\circ \pm 7.65$ ). However, the maximum KneeAngle reached is highest during SC<sub>45</sub> ( $125.16^\circ \pm 8.91$ ). Concerning the 180° representing full knee extension, the KneeAngle during SC<sub>30</sub> averaged  $35.37^\circ$ . Markolf *et al.*<sup>23</sup> mentioned that knee flexion of 40° or less increases tibial translation supported by the ACL and, consequently, the

tension sustained by the joint complex, leading to an increased risk of injury. A preventive strategy should involve increasing knee flexion to absorb impact. Cochrane *et al.*<sup>24</sup> went further by identifying angles below 30° of flexion during initial contact as a critical factor in ACL injuries during changes of directions. Consistent with Markolf *et al.*<sup>23</sup>, our results showed that for all changes of directions, the KneeAnglein exceeded 30° (SC<sub>30</sub> =  $35.37^\circ$ ; SC<sub>45</sub> =  $36.55^\circ$ ; SC<sub>45cl</sub> =  $38.57^\circ$ ) but remained below 40°, suggesting that participants subjected the joint to excessive stress, which could partially explain the previously reported increases in force.

Despite the relationship between knee flexion angle and the relative risk of joint injury, it is evident that the biomechanical strategy during direction changes is not exclusive to the knee. Several studies have attributed a determining role to other joints reflecting greater or lesser knee flexion. Imwalle *et al.*<sup>25</sup> confirmed a direct influence of the hip on the knee during direction changes. Potter *et al.*<sup>26</sup> stated that open and closed changes of directions influence the biomechanical strategies manifested by subjects, with significantly greater hip flexion in open changes than in closed changes. However, our results do not align with those reported by these authors, as hip flexion was practically the same regardless of the type of directional change. One possible explanation could be that kinematic aspects were evaluated in an agility circuit with open and closed changes of direction, where athletes performed different direction changes depending on their own technique.

Moreover, gender-based differentiation in knee kinematics has been observed, showing different biomechanical strategies related to an increased risk of knee injury between men and women<sup>27</sup>. In our study, during SC<sub>30</sub>, women exhibited a KneeAnglemax (in extension) of  $33.4^\circ$ , although without significant differences compared to men ( $p = .080$ ), who exhibited an angle close to  $36^\circ$ . Significant differences were found during SC<sub>45cl</sub>,

where women had a KneeAnglein close to 146° and men to 140°. Additionally, significant gender differences were found in KneeAnglemax ( $p = 0.006$ ), with women exhibiting more significant knee extension than men, indicating a higher injury risk. These results are consistent with those reported by James *et al.*<sup>18</sup>, who conducted a comparative study by gender during closed direction changes to 60° and found a KneeAnglein in women 5.8° lower than in men. In our study, during SC<sub>45cl</sub>, women flexed 6.4° less than men, with significant differences compared to men. This indicates that the directional change technique employed by women may be associated with an increased risk of knee injury during closed direction changes. Furthermore, in the study mentioned above, the researchers found that women significantly exceeded men in KneeAnglemax, with greater extension (64.1° and 61.4°, respectively). Our results are very similar, with men exhibiting an average of 61.8° and women 54.4°.

Regarding open changes of directions, a slightly higher, but not significant, trend in extension can be observed in women compared to men during SC<sub>30</sub>. Additionally, during SC<sub>45</sub>, the KneeAnglein values are practically the same in men and women, with an increase in knee extension in women at the moment of VGRF. Dai *et al.*<sup>28</sup> evaluated knee kinematics in men and women during SC<sub>45</sub> and also found greater extension in women compared to men at the moment of VGRF. A greater range of knee flexion during the first contact could serve as a biomechanical strategy to absorb impact and reduce the load. In this sense, women have a higher relative risk of knee injury than men, as they exhibit more significant knee extension during changes of direction<sup>29</sup>.

In conclusion, the study highlights the importance of controlling GRF and biomechanical strategies during changes in direction to reduce the risk of knee injuries. Significant gender differences in forces and accelerations during changes of directions were observed, underscoring the need to

consider specific gender factors in injury prevention. Variables such as VGRF, Force<sub>ML</sub>, and ForceAP, as well as acceleration in the vertical axis, appear to be deterministic kinetic variables that explain these gender differences and knee flexion angles. The results can be helpful in informing the design of preventive strategies in the sports field and improving the understanding of the underlying mechanisms of knee injuries during changes of direction.

### Funding

None.

### Conflict of interests

The authors declared that they have no competing interests.

### Authors' ORCID

- Adrián Feria-Madueño (AFM):  
0000-0001-7425-8694
- Timothy E. Hewett (TH):  
0000-0001-9729-7573
- Borja Sañudo (BS):  
0000-0002-9969-9573

### Contributions of authors

Conception (AFM), design and acquisition of data (AFM), formal analysis (BS), interpretation of data (AFM and BS), investigation (AFM and BS), writing the original draft preparation (AFM), writing the review and editing (TH), supervision (TH and BS).

### REFERENCES

1. Hewett TE, Myer GD, Ford KR, Heidt RS, Colosimo AJ, McLean SG, van den Bogert J, Paterno MV, Succop P. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med* 2005;33(4):492-501. doi:10.1177/0363546504269591

2. **McLean SG, Walker KB, Van Den Bogert AJ.** Effect of gender on lower extremity kinematics during rapid direction changes: An integrated analysis of three sports movements. *J Sci Med Sport* 2005;8(4):411-422. doi:10.1016/S1440-2440(05)80056-8
3. **Hewett TE, Ford KR, Myer GD.** Anterior cruciate ligament injuries in female athletes: Part 2, A meta-analysis of neuromuscular interventions aimed at injury prevention. *Am J Sports Med* 2006;34(3):490-498. doi:10.1177/0363546505282619
4. **Myer GD, Ford KR, Paterno MV, Nick TG, Hewett TE.** The effects of generalized joint laxity on risk of anterior cruciate ligament injury in young female athletes. *Am J Sports Med* 2008;36(6):1073-1080. doi:10.1177/0363546507313572
5. **Chen CH, Condello G, Chen WH, Liu TH, Tessitore A, Shiang TY, Liu C.** Effects of a novel inclined-adaptive footwear on change-of-direction performance in male athletes. *Gait Posture* 2022;94:189-194. doi:10.1016/j.gaitpost.2022.03.010
6. **Condello G, Kernozek TW, Tessitore A, Foster C.** Biomechanical analysis of a change-of-direction task in college soccer players. *Int J Sports Physiol Perform* 2016;11(1):96-101. doi:10.1123/ij-spp.2014-0458
7. **McFadden C, Daniels KAJ, Strike S.** Six methods for classifying lower-limb dominance are not associated with asymmetries during a change of direction task. *Scand J Med Sci Sports* 2022;32(1):106-115. doi:10.1111/sms.14054
8. **Olmsted LC, Carcia CR, Hertel J, Shultz SJ.** Efficacy of the star excursion balance tests in detecting reach deficits in subjects with chronic ankle instability. *J Athl Train* 2002;37(4):501-506.
9. **Maniar N, Schache AG, Pizzolato C, Opar DA.** Muscle contributions to tibiofemoral shear forces and valgus and rotational joint moments during single leg drop landing. *Scand J Med Sci Sports* 2020;30(9):1664-1674. doi:10.1111/sms.13711
10. **McBurnie AJ, Dos'Santos T, Jones PA.** Biomechanical associates of performance and knee joint loads during A 70–90° cutting maneuver in subelite soccer players. *J Strength Cond Res* 2021;35(11):3190-3198. doi:10.1519/JSC.0000000000003252
11. **Nigg BM.** The role of impact forces and foot pronation: A new paradigm. *Clin J Sport Med* 2001;11(1):2-9. doi:10.1097/00042752-200101000-00002
12. **Griffin LY, Albohm MJ, Arendt EA, Bahr R, Beynonn BD, Demaio M, Dick RW, Engbretsen L, Garrett WE Jr, Hannafin JA, Hewett TE, Huston LJ, Ireland ML, Johnson RJ, Lephart S, Mandelbaum BR, Mann BJ, Marks PH, Marshall SW, Myklebust G, Noyes FR, Powers C, Shields C Jr, Shultz SJ, Silvers H, Slauterbeck J, Taylor DC, Teitz CC, Wojtys EM, Yu B.** Understanding and preventing noncontact anterior cruciate ligament Injuries: A review of the Hunt Valley II Meeting, January 2005. *Am J Sports Med* 2006;34(9):1512-1532. doi:10.1177/0363546506286866
13. **Brughelli M, Cronin J, Levin G, Chaouachi A.** Understanding change of direction ability in sport: A review of resistance training studies. *Sports Med* 2008;38(12):1045-1063. doi:10.2165/00007256-200838120-00007
14. **Sigward SM, Powers CM.** Loading characteristics of females exhibiting excessive valgus moments during cutting. *Clin Biomech* 2007;22(7):827-833. doi:10.1016/j.clinbiomech.2007.04.003
15. **Stoffel KK, Nicholls RL, Winata AR, Dempsey AR, Boyle JJW, Lloyd DG.** Effect of ankle taping on knee and ankle joint biomechanics in sporting tasks. *Ed Sci Sports Exerc* 2010;42(11):2089-2097. doi:10.1249/MSS.0b013e3181de2e4f
16. **Wikstrom EA, Tillman MD, Chmielewski TL, Borsa PA.** Measurement and evaluation of dynamic joint stability of the knee and ankle after injury. *Sports Med* 2006;36(5):393-410. doi:10.2165/00007256-200636050-00003
17. **Liu H, Wu W, Yao W, Spang JT, Creighton RA, Garrett WE, Yu B.** Effects of knee extension constraint training on knee flexion angle and peak impact ground-reaction force. *Am J Sports Med* 2014;42(4):979-986. doi:10.1177/0363546513519323

18. James CR, Sizer PS, Starch DW, Lockhart TE, Slautebeck J. Gender differences among sagittal plane knee kinematic and ground reaction force characteristics during a rapid sprint and cut maneuver. *Res Q Exer Sport* 2004;75(1):31-38. doi:10.1080/02701367.2004.10609131
19. de Hoyo M, de la Torre A, Pradas F, Sañudo B, Carrasco L, Mateo-Cortes J, Domínguez-Cobo S, Fernandes O, Gonzalo-Skok O. Effects of eccentric overload bout on change of direction and performance in soccer players. *Int J Sports Med* 2014;36(04):308-314. doi:10.1055/s-0034-1395521
20. Chappell JD, Herman DC, Knight BS, Kirkendall DT, Garrett WE, Yu B. Effect of fatigue on knee kinetics and kinematics in stop-jump tasks. *Am J Sports Med* 2005;33(7):1022-1029. doi:10.1177/0363546504273047
21. Tillman MD, Hass CJ, Brunt D, Bennett GR. Jumping and landing techniques in elite women's volleyball. *J Sports Sci Med* 2004;3(1):30-36.
22. Pappas E, Nightingale EJ, Simic M, Ford KR, Hewett TE, Myer GD. Do exercises used in injury prevention programmes modify cutting task biomechanics? A systematic review with meta-analysis. *Br J Sports Med* 2015;49(10):673-680. doi:10.1136/bjsports-2014-093796
23. Markolf KL, O'Neill G, Jackson SR, McAllister DR. Effects of Applied quadriceps and hamstrings muscle loads on forces in the anterior and posterior cruciate ligaments. *Am J Sports Med* 2004;32(5):1144-1149. doi:10.1177/0363546503262198
24. Cochrane JL, Lloyd DG, Butfield A, Seward H, McGivern J. Characteristics of anterior cruciate ligament injuries in Australian football. *J Sci Med Sport* 2007;10(2):96-104. doi:10.1016/j.jsams.2006.05.015
25. Imwalle LE, Myer GD, Ford KR, Hewett TE. Relationship between hip and knee kinematics in athletic women during cutting maneuvers: A possible link to noncontact anterior cruciate ligament injury and prevention. *J Strength Cond Res* 2009;23(8):2223-2230. doi:10.1519/JSC.0b013e3181bc1a02
26. Potter D, Reidinger K, Szymialowicz R, Martin T, Dione D, Feinn R, Wallace D, Garbalosa JC. Sidestep and crossover lower limb kinematics during a prolonged sport-like agility test. Sidestep and crossover lower limb kinematics during a prolonged sport-like agility test. *Int J Sports Phys Ther* 2014;9(5):617-627.
27. Galloway RT, Xu Y, Hewett TE, Barber Foss K, Kiefer AW, DiCesare CA, Magnussen RA, Khoury J, Ford KR, Diekfuss JA, Grooms D, Myer GD, Montalvo AM. Age-dependent patellofemoral pain: hip and knee risk landing profiles in prepubescent and postpubescent female athletes. *Am J Sports Med* 2018;46(11):2761-2771. doi:10.1177/0363546518788343
28. Dai B, Garrett WE, Gross MT, Padua DA, Queen RM, Yu B. The Effects of 2 landing techniques on knee kinematics, kinetics, and performance during stop-jump and side-cutting tasks. *Am J Sports Med* 2015;43(2):466-474. doi:10.1177/0363546514555322
29. Mallinos A, Jones K, Davis BL. Comparison of side-cutting maneuvers versus low impact baseball swing on knee ligament loading in adolescent populations. *Clin Biomech* 2023;106:106004. doi:10.1016/j.clinbiomech.2023.106004